



Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

Sunglint filling: status update

E. Baudrez N. Clerbaux I. Decoster S. Dewitte
A. Ipe S. Nevens A. Velázquez

Royal Meteorological Institute of Belgium

23 October 2012



Outline

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

1. Sun glint filling

2. Conversion tool: HDF to netCDF

3. Regridding tool

Problem

Sunglint
filling:
status
update

RMIB


Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool


Appendices

- ▶ Under certain conditions, sun rays are reflected specularly towards the detector
- ▶ Scene identification is unreliable → radiance-to-flux conversion unreliable
- ▶ Currently there's a black patch over the area defined by the geometric condition $SGA < 15^\circ$
- ▶ Step change in flux at $SGA = 25^\circ$



- ▶ 7 July 2004, 14:30
- ▶ SW flux
- ▶ masked out over clear/cloudy ocean + over land
- ▶ (quantisation factor = 0.25)

Step change in flux at SGA = 25°



- ▶ 7 July 2004, 17:30
- ▶ SW flux
- ▶ color scale intentionally clipped from 80 to 120 W m^{-2} (clear ocean)
- ▶ step change in flux at SGA 25° transition

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

Aim

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ Estimate or restore flux in sun glint-affected area
 $SGA < 15^\circ$
- ▶ Re-estimate flux in area $15^\circ < SGA < 25^\circ$
- ▶ Fluxes applied to high-resolution ($HR = 1237 \times 1237$)
GERB and GERB-like images
- ▶ But final output should be in BARG ($= 247 \times 247$)
resolution (only?)
- ▶ Entire GERB archive to be reprocessed

Scene ID determination

Sunglint
filling:
status
update

RMIB

Sun glint
filling


Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ Over land: no problem, use as-is
- ▶ Over ocean: scene identification (cloud cover, phase & optical depth) is unreliable
- ▶ Solution: time extrapolation of scene ID just before sun glint to sun glint
 - ▶ Last valid cloud cover, phase & optical depth are retained as long as pixel is affected by sun glint
 - ▶ With a maximum of 2.5 hours (beyond that, scene ID becomes invalid)
 - ▶ Stored scene ID replaced with real scene ID as soon as available
- ▶ Not an interpolation between pre- and post-glint; it should work in near-real time

200407071430_scid_age



Radiance-to-flux conversion

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ $\text{SGA} < 15^\circ$
 - ▶ Land: radiance-to-flux conversion using up-to-date scene ID (normal procedure)
 - ▶ Clear ocean: CERES TRMM climatology (albedo + incoming solar \rightarrow flux)
 - ▶ Cloudy ocean: radiance-to-flux conversion using extrapolated scene ID
- ▶ $15^\circ < \text{SGA} < 25^\circ$
 - ▶ Land: no action required
 - ▶ Clear ocean: CERES TRMM climatology
 - ▶ Cloudy ocean: no action required

CERES TRMM climatology: flux calculation

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

CERES TRMM “flux” is calculated as follows:

$$F_{\text{TRMM}} = a \frac{F_{\text{solar}} \cos \theta}{\tilde{r}^2}$$

where

a = CERES TRMM albedo

F_{solar} = 1366 W m⁻²

θ = solar zenith angle

\tilde{r} = normalized earth–sun distance

CERES TRMM climatology: flux calculation

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

GERB flux for clear ocean is then calculated as follows:

$$F_{\text{GERB}} = \Gamma F_{\text{TRMM}}$$

where Γ (= adjustment factor) is GERB–TRMM albedo ratio:

$$\Gamma = f \left(\frac{F_{\text{GERB}}}{F_{\text{TRMM}}} \right)$$

Options for f :

1. Uniform value: $f = 1.07$
2. Time extrapolation similar to scene ID:
$$f = (F_{\text{GERB}}/F_{\text{TRMM}})_{\text{old}}$$
3. Monthly average: $f = \langle F_{\text{GERB}}/F_{\text{TRMM}} \rangle$

Example: 7 July 2004, 14:30

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



Example: 7 July 2004, 14:30

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



- ▶ flux over land restored
- ▶ reasonable-looking fill-in

Example: 7 July 2004, 14:30

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



zoom; before

Example: 7 July 2004, 14:30

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF


Regridding
tool

Appendices




- ▶ flux over land restored
- ▶ reasonable-looking fill-in
- ▶ black spot due to missing scene ID

2004_average_gerb2




average GERB-TRMM albedo
ratio
2004
average = 1.049

200407-gridded



average GERB-TRMM albedo
ratio
Jul 2004

200412-gridded



average GERB-TRMM albedo
ratio
Dec 2004

Average Γ

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

Month	GERB-2	GERB-1	Year	average
01	1.02525	0.93628	2004	1.0493
02	1.02731	0.94522	2005	1.0290
03	1.04082	0.88675	2006	1.0146
04	1.02421	0.94901	2007 (GERB-2)	1.0215
05	1.03255	0.95449	2007 (GERB-1)	0.9926
06	1.04433	0.95677	2008	0.9661
07	1.05262	0.97051	2009	0.9442
08	1.04833	0.98032	2010	0.9272
09	1.05289	0.98131	2011	0.9189
10	1.03025	0.95118		
11	1.01214	0.93965		
12	1.01207	0.92570		



Current status

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ Filling sun glint-affected areas $\text{SGA} < 15^\circ$ and $15^\circ < \text{SGA} < 25^\circ$
- ▶ Using a monthly average adjustment factor
- ▶ Available in HR and BARG resolution
- ▶ Available in HDF format as supplementary files



Next

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ Process entire archive
- ▶ Decide on a format
 - ▶ output all fields, make self-contained files
 - ▶ modified fields only, supplement existing archive files
- ▶ Output HDF files
- ▶ Twilight: model?



Outline

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

1. Sun glint filling

2. Conversion tool: HDF to netCDF

3. Regridding tool



Conversion tool: HDF to netCDF

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ A software tool to convert existing HDF files to netCDF format
- ▶ Why netCDF? Arguably easier format for end users
- ▶ netCDF libraries from version 4.0 onwards can read and write netCDF files in HDF format, but
 - ▶ incompatibility with HDF version used by GERB processing (1.6.5, netCDF requires 1.8.0)
 - ▶ HDF support for netCDF does not always come standard: many users may be stuck with older netCDF libraries
 - ▶ the HDF files would need to be tuned before they can be read by the netCDF library (netCDF won't just read any HDF file)
- ▶ Could be coupled with regridding onto latitude-longitude grid (see further)



Conversion algorithm (Prototype version)

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ (Optionally) uncompresses .hdf.gz files on the fly
- ▶ Scans the HDF file
- ▶ Creates a CDL description corresponding to the structure of the HDF file, but without the data
- ▶ Generates the netCDF file from the CDL description, with dimensions and attributes, but without the data
- ▶ Transfers the data from the HDF file to the netCDF file
- ▶ (Optionally) compress .nc files on the fly (no internal compression in netCDF classic format)
- ▶ (Currently outputs in netCDF classic format.)



Conversion tool

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ Written in Perl as a stand-alone distribution (with embedded C code)
- ▶ Could be released as open-source software
- ▶ Does not assume much about the structure of the HDF file, so could be extended to provide a general-purpose HDF-to-netCDF conversion tool



Outline

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

**Regridding
tool**

Appendices

1. Sun glint filling

2. Conversion tool: HDF to netCDF

3. Regridding tool



Regridding tool

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

- ▶ To convert data to and from
 - ▶ Sinusoidal grid
 - ▶ Regular latitude–longitude grid
 - ▶ Geostationary grid (Meteosat field of view)
- ▶ Could be used with HDF to netCDF conversion tool
- ▶ End goal: latitude–longitude projected GERB data in netCDF for end users

Regridding algorithm

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



Regridding algorithm

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



$$\blacksquare = A'_{i,j+1}$$

Regridding algorithm

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



$$A_{i,j} f_{i,j} = \boxed{A'_{i,j+1}} f_{i,j+1} + A'_{i+1,j+1} f_{i+1,j+1} + A'_{i+1,j} f_{i+1,j} + A'_{i,j} f_{i,j}$$

Regridding algorithm

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



$$A_{i,j} f_{i,j} = A'_{i,j+1} f_{i,j+1} + \boxed{A'_{i+1,j+1}} f_{i+1,j+1} + A'_{i+1,j} f_{i+1,j} + A'_{i,j} f_{i,j}$$

Regridding algorithm

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



$$A_{i,j} f_{i,j} = A'_{i,j+1} f_{i,j+1} + A'_{i+1,j+1} f_{i+1,j+1} + \boxed{A'_{i+1,j}} f_{i+1,j} + A'_{i,j} f_{i,j}$$

Regridding algorithm

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



$$A_{i,j} f_{i,j} = A'_{i,j+1} f_{i,j+1} + A'_{i+1,j+1} f_{i+1,j+1} + A'_{i+1,j} f_{i+1,j} + \boxed{A'_{i,j}} f_{i,j}$$

Acknowledgements

- ▶ Dr. Jacqueline E. Russell, Imperial College
- ▶ GERB team at RMIB
- ▶ RMIB

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

GERB-like

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



21 June 2004: **before**

- ▶ tends to increase cloud cover
- ▶ tends to decrease crispness

GERB-like

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



21 June 2004: **after**

- ▶ tends to increase cloud cover
- ▶ tends to decrease crispness



GERB-like

Sunglint
filling:
status
update


RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices



GERB-like

Sunglint
filling:
status
update

RMIB

Sun glint
filling

Conversion
tool: HDF to
netCDF

Regridding
tool

Appendices

